Workshop on Polarized Sources Targets and Polarimetry 2022 (PSTP22)



Contribution ID: 10

Type: not specified

Development of polarized sources based on molecular photodissociation

Thursday 29 September 2022 14:00 (25 minutes)

Molecular photodissociation is an innovative method for the preparation of polarized atoms and molecules. It is a fundamental chemical process that involves the absorption of one or more polarized photons by a molecule including its fragmentation into polarized atomic (or molecular) fragments. Recently, T. P. Rakitzis' group produced high densities of spin polarized hydrogen atoms applying molecular photodissociation to hydrogen halides. The obtained densities (10^{19} cm⁻³) and short production times (ns timescales) surpass by several orders of magnitude conventional methods such as spin-exchange optical pumping and Stern-Gerlach spin separation. These density and time regimes make it an ideal candidate for a broad range of applications, e.g. laser-induced acceleration from polarized gas targets and polarized five-nucleon fusion reactions (D-T, D-³He). The second has been shown to have an increased cross section by ~ 50% compared to the unpolarized case.

Category

Polarized Sources

Primary author: Mr KANNIS, Chrysovalantis (Institut für Kernphysik, Forschungszentrum Jülich, Jülich, Germany and University of Crete, Department of Physics, Herakleio, Greece)

Co-authors: Prof. BÜSCHER, Markus (Peter-Grünberg Institut, Forschungszentrum Jülich, Jülich, Germany and Institut für Laser- und Plasmaphysik, Heinrich-Heine-Universität Düsseldorf, Düsseldorf, Germany); Dr ENGELS, Ralf (Institut für Kernphysik, Forschungszentrum Jülich, Jülich, Germany); Prof. LEHRACH, Andreas (Institut für Kernphysik, Forschungszentrum Jülich, Jülich, Germany and III. Physikalisches Institut B, RWTH Aachen University, Aachen, Germany); Mr XYGKIS, Michalis (Foundation for Research and Technology Hellas, Institute of Electronic Structure and Laser, Heraklion, Crete, Greece and University of Crete, Department of Physics, Herakleio, Greece); Prof. T. PETER, Rakitzis (Foundation for Research and Technology Hellas, Institute of Laser, Heraklion, Crete, Greece and University of Crete, Department of Physics, Herakleio, Greece)

Presenter: Mr KANNIS, Chrysovalantis (Institut für Kernphysik, Forschungszentrum Jülich, Jülich, Germany and University of Crete, Department of Physics, Herakleio, Greece)

Session Classification: Polarized Sources