


A High-Field Polarized ^3He Target for Jefferson Lab's CLAS12 Spectrometer

J. Maxwell

for CLAS12 Polarized ^3He Collaboration




Polarized Sources, Targets & Polarimetry
Mainz, Germany
September 26, 2022




Outline

- 1 Polarized ^3He for Nuclear Physics
 - Optical Pumping
 - Opportunity for a New Target
- 2 Proposed Target for CLAS12
 - Design
 - Development
- 3 Outlook
 - Projected Schedule
 - Transverse Polarization?




Outline

- 1 Polarized ^3He for Nuclear Physics
 - Optical Pumping
 - Opportunity for a New Target
- 2 Proposed Target for CLAS12
 - Design
 - Development
- 3 Outlook
 - Projected Schedule
 - Transverse Polarization?




Why Polarized Helium 3?



- About 90% of the time, ^3He 's 2 proton spins are anti-aligned in a spin singlet
- ^3He spin is primarily neutron spin
- By polarizing ^3He , we have a surrogate for polarized free neutrons


Polarized ^3He Targets at JLab

- 6 GeV era: 13 experiments in Hall A
- 12 GeV era: 7 experiments approved
- Spin Exchange Optical Pumping
- 60% in-beam polarization in 10 bar gas




Metastability Exchange Optical Pumping

- 1963, Colgrove *et al* (TI)
- Pure ^3He , ~ 30 G field
- Discharge promotes states to 2^3S_1
- Laser drives polarization
- Collisions between 2^3S_1 and ground state polarize nuclei
- Requires ~ 2 mbar, > 100 K
- 10^5 faster than SEOP
- 10^4 lower pressure has limited use for scattering experiments




Metastability Exchange Optical Pumping

- 1963, Colgrove *et al* (TI)
- Pure ^3He , ~ 30 G field
- Discharge promotes states to 2^3S_1
- Laser drives polarization
- Collisions between 2^3S_1 and ground state polarize nuclei
- Requires ~ 2 mbar, > 100 K
- 10^5 faster than SEOP
- 10^4 lower pressure has limited use for scattering experiments




MEOP Double-Cell Cryo Target: Bates 88-02

- Quasi-elastic asymmetries in 1988, 1993
- MEOP pumping cell at 2 mbar, 300 K, 30 G:
40% in-beam polarization
- Cu target cell at 2 mbar, 17 k
- Cu foil beam windows ($4.6\ \mu\text{m}$)
- Cold surfaces coated with N_2 to reduce depolarization from wall interactions
- $7.2 \times 10^{32}\ ^3\text{He}/\text{cm}^2/\text{s}$ Luminosity w/ $10\ \mu\text{A}$
- P measurement performed in pumping cell
- P in target inferred from rate equations: P relaxation and diffusion




MEOP Double-Cell Cryo Target: Bates 88-02

- Quasi-elastic asymmetries in 1988, 1993
- MEOP pumping cell at 2 mbar, 300 K, 30 G:
40% in-beam polarization
- Cu target cell at 2 mbar, 17 k
- Cu foil beam windows ($4.6 \mu\text{m}$)
- Cold surfaces coated with N_2 to reduce depolarization from wall interactions
- $7.2 \times 10^{32} \text{ } ^3\text{He}/\text{cm}^2/\text{s}$ Luminosity w/ $10 \mu\text{A}$
- P measurement performed in pumping cell
- P in target inferred from rate equations: P relaxation and diffusion



High Magnetic Field MEOP

- OP not historically done at high B
 - SEOP: Increasing wall relaxation
 - MEOP: Weak hyperfine coupling...?
- Kastler-Brossel Lab at ENS in Paris found by increasing B_0 , MEOP effective at higher pressures (Nikiel-Osuchowska *et al*, Eur. Phys. J.D., 2013.)
- Near 60% at 100 mbar!
- Zeeman splitting separates states for laser pumping
 - Decouples relaxation paths
 - Creates probe peaks (Suchanek *et al.*, Euro Phys JST, 2007.)




Solid points above 1 T

From Gentile, Nacher, Saam, Walker (2017.)


High Magnetic Field MEOP

- OP not historically done at high B
 - SEOP: Increasing wall relaxation
 - MEOP: Weak hyperfine coupling...?
- Kastler-Brossel Lab at ENS in Paris found by increasing B_0 , MEOP effective at higher pressures (Nikiel-Osuchowska *et al*, Eur. Phys. J.D., 2013.)
- Near 60% at 100 mbar!
- Zeeman splitting separates states for laser pumping
 - Decouples relaxation paths
 - Creates probe peaks (Suchanek *et al.*, Euro Phys JST, 2007.)




High Magnetic Field MEOP


- OP not historically done at high B
 - SEOP: Increasing wall relaxation
 - MEOP: Weak hyperfine coupling...?
- Kastler-Brossel Lab at ENS in Paris found by increasing B_0 , MEOP effective at higher pressures (Nikiel-Osuchowska *et al*, Eur. Phys. J.D., 2013.)
- Near 60% at 100 mbar!
- Zeeman splitting separates states for laser pumping
 - Decouples relaxation paths
 - Creates probe peaks (Suchanek *et al.*, Euro Phys JST, 2007.)



Measuring Optical Pumping via Probe Peaks




Measuring Optical Pumping via Probe Peaks




High Magnetic Field MEOP for EIC

- High field MEOP techniques already being applied for nuclear physics
- BNL-MIT: Polarized ^3He Ion Source for EIC (See G. Atoian on Thursday)
- BNL's Electron Beam Ion Source operates at 5 T
- MEOP within 5 T field, transfer into EBIS for ionization and extraction
- Tests between 2 to 4 T gave nearly 90% max polarization (Maxwell *et al.*, NIM A 959, 2020)
- Installation in 2023



High Magnetic Field MEOP for EIC


- High field MEOP techniques already being applied for nuclear physics
- BNL-MIT: Polarized ^3He Ion Source for EIC (See G. Atoian on Thursday)
- BNL's Electron Beam Ion Source operates at 5 T
- MEOP within 5 T field, transfer into EBIS for ionization and extraction
- Tests between 2 to 4 T gave nearly 90% max polarization (Maxwell *et al.*, NIM A **959**, 2020)
- Installation in 2023



Results at 1.3 mbar

An Opportunity in Hall B's CLAS12

- CEBAF Large Acceptance Spectrometer for Jefferson Lab's 12 GeV upgrade
 - High luminosity electron scattering
 - Multi-particle final state response
- PR12-20-002: A program of spin-dependent electron scattering using a polarized ^3He target in CLAS12
 - P_T -dependence of n longitudinal spin structure
 - Nuclear corrections to SIDIS
 - Conditionally approved with A-rating
 - Spokespeople: Avakian, Maxwell, Milner, Nguyen
- 5 T solenoid in interaction region
- Novel target needed for standard config



Creating a New Target for CLAS12

Double-Cell Cryo Target

- Polarize at 300 K
- Transfer to 5 K target cell
- Density increase $60\times$

+


High Field MEOP

- High Polarization ($\sim 60\%$)
- High magnetic fields (5 T)
- Pressure increase $100\times$


- By combining established technologies: a new polarized target
(Maxwell, Milner, NIM A, 2021.)
- Achieve 5.4 amg, roughly half JLab SEOP target gas density
- Polarize within 5 T solenoid: CLAS12 standard configuration

Outline

- 1 Polarized ^3He for Nuclear Physics
 - Optical Pumping
 - Opportunity for a New Target
- 2 Proposed Target for CLAS12
 - Design
 - Development
- 3 Outlook
 - Projected Schedule
 - Transverse Polarization?



Proposed Target




293 K Pumping Cell

- 200 cm³ borosilicate glass
- MEOP to 60% polarization
- Annular cylindrical volume

5 K Target Cell

- 100 cm³, 20 cm long aluminum cell
- Cooled by LHe heat exchanger
- Luminosity of 2.7×10^{34} nuc/cm²/s at 0.5 μA

Proposed Target



293 K Pumping Cell


- 200 cm³ borosilicate glass
- MEOP to 60% polarization
- Annular cylindrical volume

5 K Target Cell

- 100 cm³, 20 cm long aluminum cell
- Cooled by LHe heat exchanger
- Luminosity of 2.7×10^{34} nuc/cm²/s at 0.5 μA


Depolarization Mechanisms

- Wall relaxation on Al: H_2 coatings yield days long relaxation at 4 K (Lefevre-Seguin, Low Temp. P. 1988.)
- Depolarization from transverse magnetic field gradients, dependent on pressure, temperature
 - In CLAS12 solenoid: minimal
- Beam produces $^3\text{He}_2^+$ ions: increase with density, but decrease with higher field. (Bonin, PRA, 1988)




Depolarization Mechanisms

- Wall relaxation on Al: H_2 coatings yield days long relaxation at 4 K (Lefevre-Seguin, Low Temp. P. 1988.)
- Depolarization from transverse magnetic field gradients, dependent on pressure, temperature
 - In CLAS12 solenoid: minimal
- Beam produces $^3\text{He}_2^+$ ions: increase with density, but decrease with higher field. (Bonin, PRA, 1988)




Depolarization Mechanisms

- Wall relaxation on Al: H_2 coatings yield days long relaxation at 4 K (Lefevre-Seguin, Low Temp. P. 1988.)
- Depolarization from transverse magnetic field gradients, dependent on pressure, temperature
 - In CLAS12 solenoid: minimal
- Beam produces $^3\text{He}_2^+$ ions: increase with density, but decrease with higher field. (Bonin, PRA, 1988)




Cryogenics and Heat Load

- Heat loads on 5 K target cell:
 - Beam heating (<150 mW)
 - Pumping cell at 293 K, transfer through glass and gas (<500 mW)
 - Radiative heating minimized by heat shield (<20 mW)
- Pulse-tube cryocooler for 2.5 W at 4.2 K should be sufficient (Cryomech PT425)
- JLab's Hall D cryotarget provides liquid H_2 and He_2
 - Few modifications needed to design to support a MEOP double-cell cryotarget




Cryogenics and Heat Load

- Heat loads on 5 K target cell:
 - Beam heating (<150 mW)
 - Pumping cell at 293 K, transfer through glass and gas (<500 mW)
 - Radiative heating minimized by heat shield (<20 mW)
- Pulse-tube cryocooler for 2.5 W at 4.2 K should be sufficient (Cryomech PT425)
- JLab's Hall D cryotarget provides liquid H_2 and He_2
 - Few modifications needed to design to support a MEOP double-cell cryotarget




Research Program

- High field MEOP test stand explore high field polarization vs. pressure and field, reproduce results of KBL
- Flow tests between cold and warm cells with Target Group's 4 K test stand
- Full, double-cell prototype within three years to allow in-beam tests at JLab's injector test facility




Research Program

- High field MEOP test stand explore high field polarization vs. pressure and field, reproduce results of KBL
- Flow tests between cold and warm cells with Target Group's 4 K test stand
- Full, double-cell prototype within three years to allow in-beam tests at JLab's injector test facility



Research Program

- High field MEOP test stand explore high field polarization vs. pressure and field, reproduce results of KBL
- Flow tests between cold and warm cells with Target Group's 4 K test stand
- Full, double-cell prototype within three years to allow in-beam tests at JLab's injector test facility




Beginning Polarization

- First MEOP pumping stand at JLab
 - Enclosed, interlocked system (class 1)
 - On wheels to allow transport to 5 T warm-bore solenoid
- Shakedown with low-field Helmholtz pair
 - Low-field, 668 nm liquid crystal polarimeter (Maxwell *et al.*, NIM A, 2014.)
 - Sealed ^3He cells at first
 - 60% nuclear polarization with 80 sec relaxation
- Unfortunately, pumping laser fiber sustained damage in preparation for high field tests, limiting power to 0.3 W



Beginning Polarization

- First MEOP pumping stand at JLab
 - Enclosed, interlocked system (class 1)
 - On wheels to allow transport to 5 T warm-bore solenoid
- Shakedown with low-field Helmholtz pair
 - Low-field, 668 nm liquid crystal polarimeter (Maxwell *et al.*, NIM A, 2014.)
 - Sealed ^3He cells at first
 - 60% nuclear polarization with 80 sec relaxation
- Unfortunately, pumping laser fiber sustained damage in preparation for high field tests, limiting power to 0.3 W



First High Field Tests

- High Field Polarization
 - Probe laser polarimetry (Suchanek *et al.*, Euro Phys JST, 2007.)
 - Wavelength meter
 - Python DAQ software
- Limited laser power, wavelength
 - Confirmed operation of probe polarimeter
 - Mapped pump and probe laser peaks at 2, 3, 4 T
- Despite poor laser power, saw 44% polarization




First High Field Tests

- High Field Polarization
 - Probe laser polarimetry (Suchanek *et al.*, Euro Phys JST, 2007.)
 - Wavelength meter
 - Python DAQ software
- Limited laser power, wavelength
 - Confirmed operation of probe polarimeter
 - Mapped pump and probe laser peaks at 2, 3, 4 T
- Despite poor laser power, saw 44% polarization




First High Field Tests

- High Field Polarization
 - Probe laser polarimetry (Suchanek *et al.*, Euro Phys JST, 2007.)
 - Wavelength meter
 - Python DAQ software
- Limited laser power, wavelength
 - Confirmed operation of probe polarimeter
 - Mapped pump and probe laser peaks at 2, 3, 4 T
- Despite poor laser power, saw 44% polarization




First High Field Tests

- High Field Polarization
 - Probe laser polarimetry (Suchanek *et al.*, Euro Phys JST, 2007.)
 - Wavelength meter
 - Python DAQ software
- Limited laser power, wavelength
 - Confirmed operation of probe polarimeter
 - Mapped pump and probe laser peaks at 2, 3, 4 T
- Despite poor laser power, saw 44% polarization




Preparations for Varied Pressure

- New, valved pumping cell
 - Turbopump, getter, RGA on hand
 - Gas panel in preparation
- Recondenser being installed
 - Obtaining LHe for magnet is costly, inconvenient
 - Operate for months with this setup
- Laser upgrades
 - Old laser fiber repaired
 - New laser with wider 300 GHz tuning band arrived, allows 5 T
 - Shortages meant acquisition took 1 year




Preparations for Varied Pressure

- New, valved pumping cell
 - Turbopump, getter, RGA on hand
 - Gas panel in preparation
- Recondenser being installed
 - Obtaining LHe for magnet is costly, inconvenient
 - Operate for months with this setup
- Laser upgrades
 - Old laser fiber repaired
 - New laser with wider 300 GHz tuning band arrived, allows 5 T
 - Shortages meant acquisition took 1 year



Outline

- 1 Polarized ^3He for Nuclear Physics
 - Optical Pumping
 - Opportunity for a New Target
- 2 Proposed Target for CLAS12
 - Design
 - Development
- 3 Outlook
 - Projected Schedule
 - Transverse Polarization?



Key Development Questions

- Can we reproduce the polarization performance of the KBL group?
- How much flow between cells can we produce?
- How much polarization relaxation is expected at 100 mbar and 5 K in $0.5 \mu\text{A}$ beam while inside a 5 T magnetic field?

Projected Development Schedule

- High-field polarization with sealed cells: April 2022
- Polarization at varied pressures: December 2022
- Double cryogenic cell target prototype constructed: January 2024
- Tests of full polarized target in beam: 2024

Key Development Questions

- Can we reproduce the polarization performance of the KBL group?
- How much flow between cells can we produce?
- How much polarization relaxation is expected at 100 mbar and 5 K in $0.5 \mu\text{A}$ beam while inside a 5 T magnetic field?

Projected Development Schedule

- High-field polarization with sealed cells: April 2022
- Polarization at varied pressures: December 2022
- Double cryogenic cell target prototype constructed: January 2024
- Tests of full polarized target in beam: 2024

Key Development Questions

- Can we reproduce the polarization performance of the KBL group?
- How much flow between cells can we produce?
- How much polarization relaxation is expected at 100 mbar and 5 K in $0.5 \mu\text{A}$ beam while inside a 5 T magnetic field?

Projected Development Schedule

- High-field polarization with sealed cells: April 2022
- Polarization at varied pressures: December 2022
- Double cryogenic cell target prototype constructed: January 2024
- Tests of full polarized target in beam: 2024

Key Development Questions


- Can we reproduce the polarization performance of the KBL group?
- How much flow between cells can we produce?
- How much polarization relaxation is expected at 100 mbar and 5 K in $0.5 \mu\text{A}$ beam while inside a 5 T magnetic field?

Projected Development Schedule

- High-field polarization with sealed cells: April 2022
- Polarization at varied pressures: December 2022
- Double cryogenic cell target prototype constructed: January 2024
- Tests of full polarized target in beam: 2024

Transverse Polarized ^3He in CLAS12?

- Three high-impact Hall B experiments hoped to use transversely polarized HD-Ice Target
- Transverse with CLAS12: bulk superconductor
 - Cancel 2 T \parallel CLAS12 field
 - Create a 1.5 T \perp holding field
- See G. Ciullo's talk on Tuesday
- For ^3He : same but holding field ~ 50 G
- Pumping cell in longitudinal field
- Rotate spin adiabatically in transit to target cell




MgB2 cylinder:
 86 mm \varnothing
 250 mm long
 7 mm wall

1.25 T transverse magnetization
 2.0 T axial shield
 5×10^{-3} uniformity (Y_{20}/Y_{00})
 over 20 mm radius sphere

M. Lowry

Transverse Polarized ^3He in CLAS12?


- Three high-impact Hall B experiments hoped to use transversely polarized HD-Ice Target
- Transverse with CLAS12: bulk superconductor
 - Cancel 2 T \parallel CLAS12 field
 - Create a 1.5 T \perp holding field
- See G. Ciullo's talk on Tuesday
- For ^3He : same but holding field ~ 50 G
- Pumping cell in longitudinal field
- Rotate spin adiabatically in transit to target cell



M. Lowry

Transverse Polarized ^3He in CLAS12?

- Three high-impact Hall B experiments hoped to use transversely polarized HD-Ice Target
- Transverse with CLAS12: bulk superconductor
 - Cancel 2 T \parallel CLAS12 field
 - Create a 1.5 T \perp holding field
- See G. Ciullo's talk on Tuesday
- For ^3He : same but holding field ~ 50 G
- Pumping cell in longitudinal field
- Rotate spin adiabatically in transit to target cell




Summary

- A polarized ^3He target in CLAS12 would offer a wide new class of observables of quark and gluonic structure in the neutron.
- We are developing a novel target system to operate within CLAS12 without significant modification to the spectrometer.
- Combines the successful design of the Bates 88-02 double-cell cryotarget with new developments in high magnetic field MEOP.
- Polarization tests have begun at Jefferson Lab, with an aggressive development schedule aiming to provide a target to Hall B within the next few years.

CLAS12 Polarized ^3He collaboration:

- **MIT:** X. Li, R. Milner
- **JLab:** H. Avakian, J. Brock, C. Keith, J. Maxwell, D. Meekins, D. Nguyen


Thank you for your attention!



Parameter	Bates 88-02 Target Achieved	CLAS12 Target Proposed
Pumping cell pressure (mbar)	2.6	100
Pumping cell volume (cm ³)	200	120
Target cell volume (cm ³)	79	100
Target cell length (cm)	16	20
Number of atoms in pumping cell	1.2×10^{19}	3×10^{20}
Number of atoms in target cell	6×10^{19}	1.5×10^{22}
Holding field (T)	0.003	5
Polarization	40%	60%
Incident electron beam energy (GeV)	0.574	10
Cell temperature (K)	17	5
Target thickness ($^3\text{He}/\text{cm}^2$)	1.2×10^{19}	3×10^{21}
Beam current (μA)	10	2.5
Luminosity ($^3\text{He}/\text{cm}^2/\text{s}$)	7.2×10^{32}	4.5×10^{34}

Spin-Dependent Scattering from Polarized ^3He in CLAS12

- Proposal to JLab PAC 48 for 30 days
 - P_T -dependence of the neutron longitudinal spin structure
 - Nuclear corrections to SIDIS
- Complement to studies on deuteron
- Synergistic with efforts such as SoLID
- Luminosity 2.7×10^{34} nucleons/cm 3 /s with $0.5 \mu\text{A}$ beam current
- Approved with A- rating, conditional on target development



Metastability Exchange and Spin Exchange Optical Pumping

SEOP

- Pump: alkali metals in mixture
- Transfer: spin exchange
- Low pumping rate
- Walls carefully selected
- Needs oven (473 K)
- 100 W laser typical
- **Large pressure range (1 to 13 bar)**

MEOP

- Pump: metastable population
- Transfer: metastability exchange
- High pumping rate
- Less sensitive to wall interactions
- Temperature above 100 K
- 4 W laser typical
- **Limited pressure (~ 1 mbar)**

- Pressure attainable has made SEOP the most attractive tool for JLab
- Neither have historically worked in high magnetic fields