

# Axion signatures from supernova explosions through the nucleon electric-dipole portal

*Tuesday 9 August 2022 16:43 (3 minutes)*

We consider axions coupled to nucleons and photons only through the nucleon electric-dipole moment (EDM) portal. This coupling is a model-independent feature of QCD axions, which solve the strong CP problem, and might arise as well in more general axion-like particle setups. We revise the supernova (SN) axion emission induced by the nucleon EDM coupling and refine accordingly the SN 1987A bound. Furthermore, we calculate the axion flux from a future Galactic SN and show that it might produce a peculiar and potentially detectable gamma-ray signal in a large underground neutrino detector such as the proposed Hyper-Kamiokande. The possibility to detect such a signal offers a way to search for an oscillating nucleon EDM complementary to CASPERe, without relying on the assumption that axions are a sizeable component of the dark matter. Furthermore, if axions from SN produce an observable signal, they could also lead to an amount of cosmological extra-radiation observable in future cosmic surveys.

**Primary authors:** LUCENTE, Giuseppe (Bari University & INFN Bari); Prof. MIRIZZI, Alessandro; CARENZA, Pierluca (Stockholm University, Oskar Klein Centre); Prof. GIANNOTTI, Maurizio; Dr DI LUZIO, Luca (Padua University); Mr MASTROTOTARO, Leonardo (Salerno University)

**Presenter:** LUCENTE, Giuseppe (Bari University & INFN Bari)

**Session Classification:** Poster Lightning Talks