# Fine structure of the nucleon electromagnetic form factors in the vicinity of the threshold of e+eannihilation into nucleon - antinucleon pair

#### A.I. Milstein

G.I.Budker Institute of Nuclear Physics, Novosibirsk, Russia

# Layout

- Close to the threshold of  $N\bar{N}$ , but not very close:  $e^+e^- \rightarrow p\bar{p}, n\bar{n}, \text{ mesons}; J/\psi \rightarrow p\bar{p}\pi^0(\eta); J/\psi \rightarrow p\bar{p}\rho(\omega);$  $J/\psi, \psi(2S) \rightarrow p\bar{p}\gamma; J/\psi \rightarrow \gamma\eta'\pi^+\pi^- \text{ decay.}$
- Very close to the threshold of  $N\bar{N}$ : Approach of calculation of  $\sigma_{el} (e^+e^- \to p\bar{p} \text{ and } e^+e^- \to n\bar{n})$ ,  $\sigma_{in} (e^+e^- \to \text{mesons})$ , and  $\sigma_{tot}$  cross sections.
- Our predictions for  $\sigma_{el}$ ,  $\sigma_{in}$ , and  $\sigma_{tot}$  cross sections. Discussion of isospin-violating effects.
- Conclusion





Cross section  $e^+e^- \rightarrow p\bar{p}$ ; data are from B. Aubert, et al., BaBar, Phys. Rev. D 73, 012005 (2006)

# $e^+e^- \rightarrow n\bar{n}, \ 3(\pi^+\pi^-)$ near the threshold of $N\bar{N}$ pair production



Left picture: cross section  $e^+e^- \rightarrow n\bar{n}$ ; data are from M.N. Achasov, et al., SND, Phys. Rev. D 90, 112007 (2014); right picture: cross section  $e^+e^- \rightarrow 3(\pi^+\pi^-)$ ; data are from R.R.Akhmetshin, et al., CMD3, Physics Letters, B723, 634 (2013), (black dots); B. Aubert, et al., BaBar, Phys. Rev. D 73 (2006) 052003, (green open circles)



Cross section  $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)$ ; from B. Aubert, et al., BaBar, Phys. Rev. D 73 (2006) 052003

Strong enhancement of decay probability at low invariant mass of  $p\bar{p}$  in the processes  $J/\Psi \rightarrow \gamma p\bar{p}$ ,  $B^+ \rightarrow K^+ p\bar{p}$  and  $B^0 \rightarrow D^0 p\bar{p}$ ,  $B^+ \rightarrow \pi^+ p\bar{p}$  and  $B^+ \rightarrow K^0 p\bar{p}$ ,  $\Upsilon \rightarrow \gamma p\bar{p}$ ... These effects are similar to that in  $e^+e^-$  annihilation.

One of the most natural explanation of this enhancement is final state interaction of nucleon and antinucleon

B. Kerbikov, A. Stavinsky, and V. Fedotov, Phys. Rev. C **69**, 055205 (2004); D.V. Bugg, Phys. Lett. B **598**, 8 (2004); B. S. Zou and H. C. Chiang, Phys. Rev. D **69**, 034004 (2004); B. Loiseau and S. Wycech, Phys. Rev. C **72**, 011001 (2005); A. Sibirtsev, J. Haidenbauer, S. Krewald, Ulf-G. Meiner, and A.W. Thomas, Phys. Rev. D **71**, 054010 (2005); J. Haidenbauer, Ulf-G. Meiner, A. Sibirtsev, Phys.Rev. D **74**, 017501 (2006); V.F. Dmitriev and A.I.Milstein, Phys. Lett. B 658 (2007), 13.

#### Final state interaction

Final state interaction (including annihilation channels) may be taken into account by means optical potentials:

$$V_{N\bar{N}} = U_{N\bar{N}} - iW_{N\bar{N}} \,.$$

Nijmegen, Paris, Jülich... optical potentials give the same predictions for the cross sections of elastic and inelastic scattering of unpolarized particles but essentially different predictions for spin observables! The cross section  $\sigma = \sigma_{ann} + \sigma_{cex} + \sigma_{el}$  of  $p\bar{p}$  scattering has the form

$$\sigma = \sigma_0 + (\boldsymbol{\zeta}_1 \cdot \boldsymbol{\zeta}_2) \, \sigma_1 + (\boldsymbol{\zeta}_1 \cdot \boldsymbol{\nu}) (\boldsymbol{\zeta}_2 \cdot \boldsymbol{\nu}) \, (\sigma_2 - \sigma_1) \,,$$

where  $\zeta_1$  and  $\zeta_2$  are the unit polarization vectors of the proton and antiproton, respectively.

Investigation of the process  $e^+e^- \rightarrow N\bar{N}$  gives important information for modification of optical potentials!

Near the threshold but not very close to the threshold. It is possible to neglect the proton-neutron mass difference and the Coulomb potential. Our predictions for  $e^+e^- \rightarrow N\bar{N}$  [V.F.Dmitriev, A.I.Milstein, S.G. Salnikov, PR D93, 034033 (2016)]



Left: the cross sections of  $p\bar{p}$  (red line) and  $n\bar{n}$  (green line) production, Right:  $|G_E^p/G_M^p|$  for proton. The experimental data are from J.P.Lees et al., BaBar, Phys.Rev. D 87, 092005 (2013), R.R. Akhmetshin et al., CMD3, Physics Letters B759, 634 (2016) M.N. Achasov et al.,SND, Phys. Rev. D 90, 112007 (2014).  $e^+e^- \rightarrow 6\pi$  near the threshold (via virtual  $N\bar{N}$  pair production). The cross section in the energy region between 1.7 GeV and 2.1 GeV is approximated by the formula

$$\sigma_{6\pi} = A\sigma_{\mathrm{ann}}^1 + B \cdot E + C,$$

where the best coincidence is for A = 0.56, B = 0.012 nb/MeV, C = 4.96 nb. The coefficient A agrees with the data of  $p\bar{p} \rightarrow pions$ annihilation at rest, where  $6\pi$  give ~ 55% of I = 1 contribution (C. Amsler et al., Nucl. Phys. A720, 357 (2003)).



The invariant mass spectra of  $J/\psi \rightarrow p\bar{p}\pi^0$  and  $J/\psi \rightarrow p\bar{p}\eta$  decays [V.F.Dmitriev, A.I.Milstein, S.G. Salnikov, PL B 760, 139 (2016)]:



Left:  $J/\psi \rightarrow p\bar{p}\pi^0$  decay. Right:  $J/\psi \rightarrow p\bar{p}\eta$  decay. The red band corresponds to our previous parameters of the potential and the green band corresponds to the refitted model. The phase space behavior is shown by the dashed curve.

#### $J/\psi \rightarrow p\bar{p}\gamma \ (\rho, \omega)$ decays A.I.Milstein, S.G.Salnikov, Nucl. Phys. A 966, 54 (2017)

Dominant contribution is given by the state of  $p\bar{p}$  pair with the quantum numbers  $J^{PC} = 1^{-+} ({}^{1}S_{0})$ . The invariant mass spectra in  $J/\psi \rightarrow p\bar{p}\rho(\omega)$  decays:



Left:  $J/\psi \to p\bar{p}\omega$  decay. Right:  $J/\psi \to p\bar{p}\rho$  decay.

### $J/\psi, \, \psi(2S) \to p\bar{p}\gamma \, \mathrm{decay}$

The invariant mass spectra in  $J/\psi(\psi(2S) \rightarrow p\bar{p}\gamma)$  decays:



Left:  $J/\psi \to p\bar{p}\gamma$  decay. Right:  $\psi(2S) \to p\bar{p}\gamma$  decay.



The  $\eta' \pi^+ \pi^-$  invariant mass spectrum in  $J/\psi \rightarrow \gamma \eta' \pi^+ \pi^-$  decay:

The thin line shows the contribution of non-NN channels. Vertical dashed line is the  $N\bar{N}$  threshold.

### Very close to the thresholds.

### (A.I.Milstein, S.G.Salnikov, arXiv:1804.01283)

It is necessary to take also into account the proton-neutron mass difference and the Coulomb potential. The coupled-channels radial Schrdinger equation for the  ${}^{3}S_{1} - {}^{3}D_{1}$  states reads

$$\begin{bmatrix} p_r^2 + \mu \mathcal{V} - \mathcal{K}^2 \end{bmatrix} \Psi = 0, \qquad \Psi^T = (u^p, w^p, u^n, w^n),$$
$$\mathcal{K}^2 = \begin{pmatrix} k_p^2 \mathbb{I} & 0\\ 0 & k_n^2 \mathbb{I} \end{pmatrix}, \qquad \mathbb{I} = \begin{pmatrix} 1 & 0\\ 0 & 1 \end{pmatrix}, \qquad \mu = \frac{1}{2} \left( m_p + m_n \right)$$
$$k_p^2 = \mu E, \qquad k_n^2 = \mu (E - 2\Delta), \qquad \Delta = m_n - m_p,$$

where  $(-p_r^2)$  is the radial part of the Laplace operator,  $u^p(r)$ ,  $w^p(r)$  and  $u^n(r)$ ,  $w^n(r)$  are the radial wave functions of a protonantiproton or neutron-antineutron pair with the orbital angular momenta L = 0 and L = 2, respectively,  $m_p$  and  $m_n$  are the proton and neutron masses, E is the energy of a system counted from the  $p\bar{p}$ threshold.

#### The optical potential.

 $\mathcal{V}$  is the matrix  $4 \times 4$  which accounts for the  $p\bar{p}$  interaction and  $n\bar{n}$  interaction as well as a transition  $p\bar{p} \leftrightarrow n\bar{n}$ . This matrix can be written in a block form as

$$\mathcal{V} = \begin{pmatrix} \mathcal{V}^{pp} & \mathcal{V}^{pn} \\ \mathcal{V}^{pn} & \mathcal{V}^{nn} \end{pmatrix},$$

where the matrix elements read

$$\begin{split} \mathcal{V}^{pp} &= \frac{1}{2} (\mathcal{U}^1 + \mathcal{U}^0) - \frac{\alpha}{r} \mathbb{I} + \mathcal{U}_{cf} \,, \qquad \mathcal{V}^{nn} = \frac{1}{2} (\mathcal{U}^1 + \mathcal{U}^0) + \mathcal{U}_{cf} \,, \\ \mathcal{V}^{pn} &= \frac{1}{2} (\mathcal{U}^0 - \mathcal{U}^1) \,, \qquad \mathcal{U}_{cf} = \frac{6}{\mu r^2} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \,, \\ \mathcal{U}^I &= \begin{pmatrix} V_S^I & -2\sqrt{2} V_T^I \\ -2\sqrt{2} V_T^I & V_D^I - 2V_T^I \end{pmatrix} \,. \end{split}$$

 $V_S^I(r)$ ,  $V_D^I(r)$ , and  $V_T^I(r)$  are the terms in the potential  $V^I$  of the strong NN interaction, corresponding to the isospin I,

$$V^{I} = V_{S}^{I}(r)\delta_{L0} + V_{D}^{I}(r)\delta_{L2} + V_{T}^{I}(r)\left[6\left(\boldsymbol{S}\cdot\boldsymbol{n}\right)^{2} - 4\right]$$

Here S is the spin operator of the produced pair (S = 1) and n = r/r. The optical potential V is expressed via the potentials  $\tilde{U}^I$  as follows

$$V(r) = \widetilde{U}^0 + (\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2) \, \widetilde{U}^1,$$

 $au_{1,2}$  are the isospin Pauli matrices. The terms  $V^I_{S,D,T}$  are

 $V_i^1(r) = \widetilde{U}_i^0(r) + \widetilde{U}_i^1(r) \,, \ V_i^0(r) = \widetilde{U}_i^0(r) - 3\widetilde{U}_i^1(r) \,, \ i = S, D, T \,.$ 

The potentials  $\widetilde{U}_i^I(r)$  consist of the real and imaginary parts:

$$\begin{split} \widetilde{U}_i^0(r) &= \left( U_i^0 - i \, W_i^0 \right) \theta \left( a_i^0 - r \right), \\ \widetilde{U}_i^1(r) &= \left( U_i^1 - i \, W_i^1 \right) \theta \left( a_i^1 - r \right) + U_i^{\pi}(r) \theta \left( r - a_i^1 \right), \end{split}$$

where  $\theta(x)$  is the Heaviside function,  $U_i^I$ ,  $W_i^I$ ,  $a_i^I$  are free parameters fixed by fitting the experimental data, and  $U_i^{\pi}(r)$  are the terms in the pion-exchange potential.

	$\widetilde{U}^0_S$	$\widetilde{U}_D^0$	$\widetilde{U}_T^0$	$\widetilde{U}_S^1$	$\widetilde{U}_D^1$	$\widetilde{U}_T^1$
$U_i({ m MeV})$	$-458^{+10}_{-12}$	$-184^{+17}_{-20}$	$-43^{+4}_{-3}$	$1.9\pm0.6$	$991^{+13}_{-15}$	$-4.5\substack{+0.2\-0.1}$
$W_i({ m MeV})$	$247\pm5$	$82^{+13}_{-7}$	$-31^{+2}_{-6}$	$-8.9\substack{+0.8\\-0.5}$	$\mathbf{5^{+14}_{-20}}$	$1.7\substack{+0.2\-0.1}$
$a_i({ m fm})$	$0.531\substack{+0.007\\-0.006}$	$1.17\substack{+0.02\\-0.03}$	$0.74\pm0.03$	$1.88\pm0.02$	$0.479 \pm 0.003$	$2.22\pm0.03$
g	$g_p = 0.338 \pm 0.004$			$g_n = -0.15 - 0.33i \pm 0.01$		

The parameters of the short-range potential.

The asymptotic forms of four independent regular solutions Solutions, which have no singularities at r = 0, at large distances are

$$\Psi_{1R}^{T}(r) = \frac{1}{2i} \left( S_{11}\chi_{p0}^{+} - \chi_{p0}^{-}, S_{12}\chi_{p2}^{+}, S_{13}\chi_{n0}^{+}, S_{14}\chi_{n2}^{+} \right),$$
  

$$\Psi_{2R}^{T}(r) = \frac{1}{2i} \left( S_{21}\chi_{p0}^{+}, S_{22}\chi_{p2}^{+} - \chi_{p2}^{-}, S_{23}\chi_{n0}^{+}, S_{24}\chi_{n2}^{+} \right),$$
  

$$\Psi_{3R}^{T}(r) = \frac{1}{2i} \left( S_{31}\chi_{p0}^{+}, S_{32}\chi_{p2}^{+}, S_{33}\chi_{n0}^{+} - \chi_{n0}^{-}, S_{34}\chi_{n2}^{+} \right),$$
  

$$\Psi_{4R}^{T}(r) = \frac{1}{2i} \left( S_{41}\chi_{p0}^{+}, S_{42}\chi_{p2}^{+}, S_{43}\chi_{n0}^{+}, S_{44}\chi_{n2}^{+} - \chi_{n2}^{-} \right).$$

Here  $S_{ij}$  are some functions of the energy and

$$\chi_{pl}^{\pm} = \frac{1}{k_p r} \exp\left[\pm i \left(k_p r - l\pi/2 + \eta \ln(2k_p r) + \sigma_l\right)\right],$$
  
$$\chi_{nl}^{\pm} = \frac{1}{k_n r} \exp\left[\pm i \left(k_n r - l\pi/2\right)\right],$$
  
$$\sigma_l = \frac{i}{2} \ln \frac{\Gamma\left(1 + l + i\eta\right)}{\Gamma\left(1 + l - i\eta\right)}, \qquad \eta = \frac{m_p \alpha}{2k_p},$$

where  $\Gamma(x)$  is the Euler  $\Gamma$  function.

#### The amplitude of $e^+e^- \rightarrow N\bar{N}$ near the threshold

In the non-relativistic approximation the amplitudes in units  $\pi \alpha / \mu^2$  are

$$T_{\lambda'\lambda}^{p\bar{p}} = \sqrt{2}G_s^p(\boldsymbol{e}_{\lambda'}\cdot\boldsymbol{\epsilon}_{\lambda}^*) + G_d^p\left[(\boldsymbol{e}_{\lambda'}\cdot\boldsymbol{\epsilon}_{\lambda}^*) - 3(\hat{\boldsymbol{k}}\cdot\boldsymbol{e}_{\lambda'})(\hat{\boldsymbol{k}}\cdot\boldsymbol{\epsilon}_{\lambda}^*)\right],$$
  
$$T_{\lambda'\lambda}^{n\bar{n}} = \sqrt{2}G_s^n(\boldsymbol{e}_{\lambda'}\cdot\boldsymbol{\epsilon}_{\lambda}^*) + G_d^n\left[(\boldsymbol{e}_{\lambda'}\cdot\boldsymbol{\epsilon}_{\lambda}^*) - 3(\hat{\boldsymbol{k}}\cdot\boldsymbol{e}_{\lambda'})(\hat{\boldsymbol{k}}\cdot\boldsymbol{\epsilon}_{\lambda}^*)\right],$$

where  $e_{\lambda'}$  is a virtual photon polarization vector, corresponding to the spin projection  $J_z = \lambda' = \pm 1$ ,  $\epsilon_{\lambda}$  is the spin-1 function of  $N\bar{N}$ pair,  $\lambda = \pm 1$ , 0 is the spin projection on the nucleon momentum k, and  $\hat{k} = k/k$ . In the non-relativistic approximation the standard formula for the differential cross section of  $N\bar{N}$  pair production in  $e^+e^-$  annihilation reads

$$\frac{d\sigma^N}{d\Omega} = \frac{k_N \alpha^2}{16\mu^3} \left[ \left| G_M^N(E) \right|^2 \left( 1 + \cos^2 \theta \right) + \left| G_E^N(E) \right|^2 \sin^2 \theta \right].$$

Here  $\theta$  is the angle between the electron (positron) momentum and the momentum of the final particle.

The proton and neutron Sachs form factors are:

$$G_{M}^{p} = G_{s}^{p} + \frac{1}{\sqrt{2}}G_{d}^{p}, \quad G_{E}^{p} = G_{s}^{p} - \sqrt{2}G_{d}^{p},$$
$$G_{M}^{n} = G_{s}^{n} + \frac{1}{\sqrt{2}}G_{s}^{n}, \quad G_{E}^{n} = G_{s}^{n} - \sqrt{2}G_{d}^{n}.$$

Thus,

$$G_E^N/G_M^N = (1-\sqrt{2}\,x)/(1+x/\sqrt{2}), \quad x = G_d^N/G_s^N\,.$$

The explicit forms of the form factors are

$$\begin{split} G_s^p &= g_p u_{1R}^p(0) + g_n u_{1R}^n(0), \quad G_d^p = g_p u_{2R}^p(0) + g_n u_{2R}^n(0), \\ G_s^n &= g_p u_{3R}^p(0) + g_n u_{3R}^n(0), \quad G_d^n = g_p u_{4R}^p(0) + g_n u_{4R}^n(0), \end{split}$$

The quantities  $u_{iR}^p(r)$  and  $u_{iR}^n(r)$  denote the first and third components of the regular solutions  $\Psi_{iR}(r)$ . The amplitudes  $g_p$  and  $g_n$  can be considered as the energy independent parameters.

# The elastic $N\bar{N}$ pair production cross section.

The integrated cross sections of the nucleon-antinucleon pair production have the form

$$\sigma_{\rm el}^p = \frac{\pi k_p \alpha^2}{4\mu^3} \left[ |G_s^p|^2 + |G_d^p|^2 \right], \quad \sigma_{\rm el}^n = \frac{\pi k_n \alpha^2}{4\mu^3} \left[ |G_s^n|^2 + |G_d^n|^2 \right].$$

The label "el" indicates that the process is elastic, i.e., a virtual  $N\overline{N}$  pair transfers to a real pair in a final state.

The inelastic cross section  $\sigma_{in}$ .

There is also an inelastic process when a virtual NN pair transfers into mesons in a final state. The total cross section  $\sigma_{tot}$ , is

$$\sigma_{\rm tot} = \sigma_{\rm el}^p + \sigma_{\rm el}^n + \sigma_{\rm in} \,. \tag{1}$$

The total cross section may be expressed via the Green's function  $\mathcal{D}(r, r'|E)$  of the wave equation

$$\sigma_{\text{tot}} = \frac{\pi \alpha^2}{4\mu^3} \text{Im} \Big[ \mathcal{G}^{\dagger} \mathcal{D} \left( 0, \ 0 | E \right) \mathcal{G} \Big], \qquad \mathcal{G}^T = \left( g_p, \ 0, \ g_n, \ 0 \right), \qquad (2)$$

where the function  $\mathcal{D}(r, r'|E)$  satisfies the equation

$$\left[p_r^2 + \mu \mathcal{V} - \mathcal{K}^2\right] \mathcal{D}\left(r, \, r' | E\right) = \frac{1}{rr'} \delta\left(r - r'\right) \,. \tag{3}$$

The function  $\mathcal{D}(r, 0|E)$  can be written as

$$\mathcal{D}(r, 0|E) = k_p \left[ \Psi_{1N}(r) \Psi_{1R}^T(0) + \Psi_{2N}(r) \Psi_{2R}^T(0) \right] + k_n \left[ \Psi_{3N}(r) \Psi_{3R}^T(0) + \Psi_{4N}(r) \Psi_{4R}^T(0) \right],$$

Non-regular solutions are defined by their asymptotic behavior at large distances:

$$u_{1N}^{p}(r) = \chi_{p0}^{+}, \quad w_{2N}^{p}(r) = \chi_{p2}^{+}, \quad u_{3N}^{n}(r) = \chi_{n0}^{+}, \quad w_{4N}^{n}(r) = \chi_{n2}^{+}.$$
(4)

All other elements  $\psi_i$  of the non-regular solutions satisfy the relation

$$\lim_{r \to \infty} r \psi_i(r) = 0 \,.$$

# Results.



Comparison of our predictions with the data for  $\sigma_{el}$  of  $e^+e^- \rightarrow p\bar{p}$ The experimental data are from J.P.Lees et al., BaBar, Phys.Rev. D 87, 092005 (2013).



 $\sigma_{el}$  for  $p\bar{p}$  (left) and  $n\bar{n}$  (right) as a function of E of a pair. Solid curves are the exact results, dashed curves are obtained at  $\Delta = 0$  and without account for the Coulomb potential, dotted curve in the left picture is obtained at  $\Delta = 0$  and with account for the Coulomb potential, dashdotted curve in the left picture corresponds to the approximation

$$\sigma_{\rm el} = C \sigma_{\rm el}^{(0)}, \quad C = \frac{2\pi\eta}{1 - e^{-2\pi\eta}}, \quad \eta = \frac{m_p \alpha}{2k_p}$$



 $\sigma_{\text{tot}}$  (left) and  $\sigma_{\text{in}}$  (right) as a function of E. Solid curves correspond to the exact results, dashed curves are the results, obtained at  $\Delta = 0$ and without account for the Coulomb interaction, dotted curves are obtained at  $\Delta = 0$  and with account for the Coulomb potential, and dash-dotted curves are obtained at  $\Delta \neq 0$  and without account for the Coulomb potential. Vertical lines show the thresholds of  $p\bar{p}$  and  $n\bar{n}$ pair production.



The cross sections  $\sigma_{tot}$ ,  $\sigma_{in}$ ,  $\sigma_{el}^p$ , and  $\sigma_{el}^n$  as a function of E.



Wave functions at origin for  $p\bar{p}$  in a final state. Solid curves are the exact results, dashed curves are the results, obtained without account for the Coulomb interaction.



Wave functions at origin for  $n\bar{n}$  in a final state. The Coulomb interaction is unimportant.

# Conclusion

- We have investigated in detail the energy dependence of the cross sections of  $p\bar{p}$ ,  $n\bar{n}$ , and meson production in  $e^+e^-$  annihilation in the vicinity of the  $p\bar{p}$  and  $n\bar{n}$  thresholds.
- Unusual phenomena are related to the interaction at large distances ("nuclear physics" of elementary particles).
- An importance of the isospin-violating effects (proton-neutron mass difference and the Coulomb interaction) is elucidated.
- Commonly accepted factorization approach for the account of the Coulomb potential does not work well enough in the vicinity of the thresholds.
- The results of SND and CMD-3 obtained at  $e^+e^-$  collider VEPP-2000 will give an important contribution to understanding of the phenomena.