

Pole determination of first discovered pentaquark with strangeness

Phys. Rev. D 108, L011501 (2023)


Satoshi Nakamura (Shandong Univ.)

Collaborator: Jia-Jun Wu (Univ. of Chinese Academy of Sciences)

Introduction

LHCb data on $B^- \rightarrow J/\psi \Lambda \bar{p}$

PRL 131, 031901 (2023)



Discovery ($>10\sigma$) of first pentaquark candidates with strangeness ($c\bar{c}uds$)

$P_{\psi s}^{\Lambda}(4338)$ properties:

$$M = 4338.2 \pm 0.7 \pm 0.4 \text{ MeV} \quad (\text{mass})$$

$$\Gamma = 7.0 \pm 1.2 \pm 1.3 \text{ MeV} \quad (\text{width})$$


$$J^P = 1/2^- \quad (\text{spin parity})$$

M , Γ , and J^P are crucial information to understand

the nature (hadron molecule, compact pentaquark, etc.) of $P_{\psi s}^{\Lambda}(4338)$

Q: M and Γ determined by LHCb are reliable ?

Basic assumption in LHCb amplitude analysis : Breit-Wigner (BW) amplitude well simulates $P_{\psi_s}^\Lambda(4338)$



BW fit

Resonance-like peak is right on the $\Xi_c \bar{D}$ threshold

→ BW fit (no unitarity) ignores important physics

- Resonance-like $\Xi_c \bar{D}$ threshold cusp appears (kinematical effect) even without a pole

In the presence of a pole

- Distortion of peak shape due to $\Xi_c \bar{D}$ branch point and cut
- Rapid increase of width just above $\Xi_c \bar{D}$ threshold

M and Γ from BW fit are questionable

What needs to be done ?


Ans. Replace BW fit with the proper pole extraction method ← The main task of this work

- Unitary coupled-channel amplitude is fitted to data
- Poles on relevant Riemann sheets are searched by analytic continuation of the amplitude



The pole value is:

- Important knowledge reflecting QCD dynamics
- Primary basis to study the nature of the pentaquark

Possible $P_{\psi_s}^\Lambda(4255)$?



Possible $P_{\psi_s}^\Lambda(4255)$?



Possibility : $\Lambda_c \bar{D}_s$ threshold cusp is enhanced by a nearby pole $P_{\psi_s}^\Lambda(4255)$ → to be examined

In this work

Conduct amplitude analysis on the LHCb data for $B^- \rightarrow J/\psi \Lambda \bar{p}$



$M_{J/\psi \Lambda}$, $M_{J/\psi \bar{p}}$, $M_{\Lambda \bar{p}}$, and $\cos \theta_{K^*}$ distribution data are simultaneously fitted with a model in which $\Xi_c \bar{D} - \Lambda_c \bar{D}_s$ coupled-channel amplitude is implemented

Based on the $\Xi_c \bar{D} - \Lambda_c \bar{D}_s$ amplitude, we address:

- (i) Pole position of $P_{\psi_s}^\Lambda(4338)$
- (ii) Possibility that $P_{\psi_s}^\Lambda(4338)$ is merely a threshold cusp (no pole)
- (iii) Implication of large fluctuation at $\Lambda_c \bar{D}_s$ threshold






Model


Model for $B^- \rightarrow J/\psi \Lambda \bar{p}$




All visible structures are at thresholds

→ threshold cusps enhanced or suppressed by hadron scattering and pole (reasonable assumption)






$\Lambda_c \bar{D}_s - \Xi_c \bar{D} - J/\psi \Lambda$ (T) and $\bar{\Lambda}_c D - J/\psi \bar{p}$ (T_2) coupled-channel s -wave amplitudes are implemented



Data-driven MB contact interactions (V) and coupled-channel unitarity : idea similar to K -matrix approach

Transitions to $J/\psi \Lambda$ and $J/\psi \bar{p}$ channels are treated perturbatively; heavy-quark exchange is expected to be weak





cf. LHCb used non-resonant p -wave $J/\psi \bar{p}$ pair production \rightarrow 84% fit fraction [counter intuitive p -wave dominance]

Results

Dalitz plot for $B^- \rightarrow J/\psi \Lambda \bar{p}$

LHCb seminar
07/05/2022







Note: No smearing due to experimental resolution is applied
→ Peak structures seem sharper than data

Fit to LHCb data for $B^- \rightarrow J/\psi \Lambda \bar{p}$

Four distribution data
are simultaneously fitted

Smearing with resolution
integration in each bin



→ histogram





$\cos \theta_{K^*} \equiv \hat{p}_{\Lambda} \cdot \hat{p}_{J/\psi}$
in $\Lambda \bar{p}$ CM frame

$\chi^2/\text{ndf} \sim 1.21$
9 parameters



Fit to LHCb data




Repulsive $\bar{\Lambda}_c D$ interaction causes suppressed cusp, increasing $M_{J/\psi\bar{p}}$ lineshape








Contribution of phase-space-like shape







$\Lambda_c \bar{D}_s - \Xi_c \bar{D}$ coupled-channel scattering causes poles near $\Lambda_c \bar{D}_s$ and $\Xi_c \bar{D}$ thresholds
 \rightarrow enhanced threshold cusps







Pole effects on the physical energy region (spectrum lineshape) are significantly screened by branch cut

Resonance-like lineshapes are caused by kinematical threshold cusps, and poles moderately enhance them

Poles are from $\Xi_c \bar{D} - \Lambda_c \bar{D}_s$ s -wave amplitude $\rightarrow J^P = 1/2^-$ poles; consistent with LHCb analysis result

Without
coupled-channel
effects


$\Xi_c \bar{D} \rightarrow \Xi_c \bar{D}$ interaction only

$\Xi_c \bar{D}$ bound state \rightarrow

$\Lambda_c \bar{D}_s \rightarrow \Lambda_c \bar{D}_s$ interaction only

$\Lambda_c^+ D_s^-$ virtual state \rightarrow

- (A) $P_{\psi_s}^\Lambda(4255)$ pole doesn't exist; the fluctuation is just statistical
- (B) $\Xi_c \bar{D} \rightarrow \Xi_c \bar{D}$ interaction has energy dependence (default result is from energy-independent interaction)
- (C) Nearby poles do not exist; peak structures in data are solely from threshold cusps




(A) and (B) have fit quality comparable to default fit
 $\Lambda_c \bar{D}_s$ threshold cusp w/o pole

(C) fit in $P_{\psi_s}^\Lambda(4338)$ peak region is visibly worse

→ $P_{\psi_s}^\Lambda(4338)$ is not merely a threshold cusp
 a nearby pole exists

- (A) $P_{\psi_s}^\Lambda(4255)$ pole doesn't exist; the fluctuation is just statistical
- (B) $\Xi_c \bar{D} \rightarrow \Xi_c \bar{D}$ interaction has energy dependence (default result is from energy-independent interaction)
- (C) Nearby poles do not exist; peak structures in data are solely from threshold cusps



(A) and (B) have fit quality comparable to default fit
 $\Lambda_c \bar{D}_s$ threshold cusp w/o pole

(C) fit in $P_{\psi_s}^\Lambda(4338)$ peak region is visibly worse

→ $P_{\psi_s}^\Lambda(4338)$ is not merely a threshold cusp
 a nearby pole exists

$P_{\psi_S}^\Lambda(4338)$

Pole locations for other solutions

Solution	E_{pole} (MeV)	sheet ($s_{\Lambda_c \bar{D}_s}$ $s_{\Xi_c^0 \bar{D}^0}$ $s_{\Xi_c^+ D^-}$)	w/o coupled-channel
(default)	$(4338.2 \pm 1.4) - (1.9 \pm 0.5)i$	(<i>upp</i>)	$\Xi_c \bar{D}$ bound pole
(A)	$(4331.9 \pm 4.2) + (5.6 \pm 6.4)i$	(<i>ppu</i>) + (<i>pup</i>), (<i>puu</i>) poles	$\Xi_c \bar{D}$ virtual pole
(B)	$(4338.0 \pm 4.1) - (6.2 \pm 7.3)i$	(<i>uuu</i>) + (<i>upp</i>), (<i>uup</i>) poles	$\Xi_c \bar{D}$ resonance pole

$P_{\psi_S}^\Lambda(4255)$

(default)	4254.7 ± 0.4	(<i>upp</i>)	$\Lambda_c \bar{D}_s$ virtual pole
-----------	------------------	----------------	------------------------------------

Depending on the solutions, $P_{\psi_S}^\Lambda(4338)$ pole is located on different Riemann-sheet \rightarrow More data needed

- Higher statistics $B^- \rightarrow J/\psi \Lambda \bar{p}$ not only pin down existence of $P_{\psi_S}^\Lambda(4255)$ but constrain $P_{\psi_S}^\Lambda(4338)$ pole sheet
- $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ should show pole effect more clearly, since no shrinking phase-space near kinematical end
 \rightarrow favor or disfavor resonance pole (larger width)

Summary

Summary

- Amplitude analysis of new LHCb data of $B^- \rightarrow J/\psi \Lambda \bar{p}$
- $M_{J/\psi \Lambda}$, $M_{J/\psi \bar{p}}$, $M_{\Lambda \bar{p}}$, and $\cos \theta_{K^*}$ distributions are fitted simultaneously; $\chi^2/\text{ndf} \sim 1.21$
- First pole determination of first discovered pentaquark candidate with strangeness $P_{\psi_s}^\Lambda(4338)$
 - important in its own right, knowledge of QCD dynamics
 - primary basis to study the nature of $P_{\psi_s}^\Lambda(4338)$
- Data disfavors hypothesis that the $P_{\psi_s}^\Lambda(4338)$ peak is just a kinematical effect
- $P_{\psi_s}^\Lambda(4255)$ might exist, and its pole is determined
- Alternative solutions have $P_{\psi_s}^\Lambda(4338)$ poles on different Riemann sheets
 - future data needed to discriminate them

Recent theoretical papers identified their $\Xi_c \bar{D}$ bound states with $P_{\psi_s}^\Lambda(4338)$

Common argument : their $\Xi_c \bar{D}$ bound state energy is consistent with M and Γ from LHCb analysis

$$M = \underline{4338.2 \pm 0.7 \pm 0.4 \text{ MeV}} \quad \Gamma = 7.0 \pm 1.2 \pm 1.3 \text{ MeV} \quad (\text{LHCb})$$

1.0 (2.9) MeV above $\Xi_c^+ D^-$ ($\Xi_c^0 \bar{D}^0$) threshold, indicating resonance not bound state, even considering error

→ The LHCb result rules out (or disfavors) the bound state solutions

Good news for $\Xi_c \bar{D}$ bound state model

BW fit employed in the LHCb analysis is unsuitable to describe $P_{\psi_s}^\Lambda(4338)$

Our proper pole extraction (default model) supports $\Xi_c \bar{D}$ bound state solution for $P_{\psi_s}^\Lambda(4338)$


Theoretical calculations of $P_{\psi_s}^\Lambda(4338)$ should be compared with our pole values; not BW values

Backup

Solution		E_{pole} (MeV)	sheet ($s_{\Lambda_c \bar{D}_s}$ $s_{\Xi_c^0 \bar{D}^0}$ $s_{\Xi_c^+ D^-}$)	
default	$P_{\psi_s}^\Lambda(4338)$	$(4338.0 \pm 1.1) - (1.7 \pm 0.4) i$	(upp)	$\Xi_c \bar{D}$ bound pole
	$P_{\psi_s}^\Lambda(4255)$	4254.6 ± 0.5	(upp)	$\Lambda_c \bar{D}_s$ virtual pole
(A)	$P_{\psi_s}^\Lambda(4338)$	$(4334.2 \pm 3.6) + (5.3 \pm 5.7) i$	(ppu)	
		$(4330.7 \pm 4.0) + (3.9 \pm 5.4) i$	(pup)	
		$(4336.4 \pm 1.4) - (0.1 \pm 1.3) i$	(upu)	
(B)	$P_{\psi_s}^\Lambda(4338)$	$(4338.9 \pm 1.7) - (2.2 \pm 0.7) i$	(upp)	$\Xi_c \bar{D}$ resonance pole
		$(4338.8 \pm 1.9) - (4.3 \pm 2.1) i$	(uup)	
		$(4337.3 \pm 1.3) - (5.1 \pm 2.5) i$	(uuu)	


Impact of pole on amplitude on the physical energy axis (data)

- Pole far from threshold



Breit-Wigner form is
good approximation

- Pole near threshold



- The existence of branch point (threshold) distorts the spectrum shape (data)
- Breit-Wigner ignores branch point (no unitarity)
→ not suitable for pole near threshold

This physical energy region is most affected by pole